

1. A method for preventing carbon tetrachloride induced liver damage in mammals,
said method comprising the steps of:

administering, to a patient, a formulation comprising processed *Morinda*
Citrifolia.

5

2. The method of claim 1, wherein said *Morinda Citrifolia* is administered in
dosages comprising at least two ounces of processed *Morinda Citrifolia* juice twice daily
on an empty stomach for a period of at least two months.

TOP SECRET - DEFENSE

10 3. The method of claim 1, wherein said *Morinda Citrifolia* is in juice form.

4. The method of claim 1, wherein said *Morinda Citrifolia* is in solid form.

5. The method of claim 1, wherein said *Morinda Citrifolia* is in powder form.

15

6. The method of claim 1, wherein

7. The method of claim 1, wherein said formulation comprises said *Morinda*
Citrifolia in an amount between about 0.1 and 5 percent by weight, said formulation
20 being administered for a period of at least seven days.

8. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 5 and 10 percent by weight, said formulation being administered for a period of at least seven days.

5 9. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 10 and 15 percent by weight, said formulation being administered for a period of at least seven days.

10. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 15 and 20 percent by weight, said formulation being administered for a period of at least seven days.

11. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 20 and 25 percent by weight, said formulation being administered for a period of at least seven days.

12. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 25 and 30 percent by weight, said formulation being administered for a period of at least seven days.

20 13. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 30 and 35 percent by weight, said formulation being administered for a period of at least seven days.

14. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 35 and 40 percent by weight, said formulation being administered for a period of at least seven days.

5

15. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 40 and 45 percent by weight, said formulation being administered for a period of at least seven days.

10 16. The method of claim 1, wherein said formulation comprises said *Morinda*
Citrifolia in an amount between about 45 and 50 percent by weight, said formulation
being administered for a period of at least seven days.

17. The method of claim 1, wherein said formulation comprises said *Morinda*
15 *Citrifolia* in an amount between about 50 and 55 percent by weight, said formulation
being administered for a period of at least seven days.

18. The method of claim 1, wherein said formulation comprises said *Morinda*
Citrifolia in an amount between about 55 and 60 percent by weight, said formulation
20 being administered for a period of at least seven days.

19. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 60 and 65 percent by weight, said formulation being administered for a period of at least seven days.

Cont'd

5 20. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 65 and 70 percent by weight, said formulation being administered for a period of at least seven days.

10 21. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 70 and 75 percent by weight, said formulation being administered for a period of at least seven days.

15 22. The method of claim 1, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 75 and 80 percent by weight, said formulation being administered for a period of at least seven days.

23. A method of inhibiting further liver damage in mammals, said method comprising the steps of:

administering, to a patient, a formulation comprising processed *Morinda Citrifolia*.

5

24. The method of claim 23, wherein said formulation comprises ten percent said *Morinda Citrifolia*, said formulation being administered for a period of at least seven days.

10 25. The method of claim 23, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 0.1 and 80 percent by weight, said formulation being administered for a period of at least seven days.

26. A method for preventing cancerous growth, in the liver of mammals, at the initiation stages of carcinogenesis by blockage of carcinogen-DNA adduct formation, said method comprising the steps of:

administering, to a patient, a formulation comprising processed *Morinda*

5 *Citrifolia*.

27. The method of claim 26, wherein said formulation comprises said *Morinda Citrifolia* in an amount between about 0.1 and 80 percent by weight, said formulation being administered for a period of at least seven days.

10

Add A2
Add B2